




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 307/79, 333/54, A61K 31/34	A1	(11) International Publication Number: WO 97/28143 (43) International Publication Date: 7 August 1997 (07.08.97)
(21) International Application Number: PCT/US97/01308 (22) International Filing Date: 30 January 1997 (30.01.97) (30) Priority Data: 08/595,158 1 February 1996 (01.02.96) US (71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventor: SCHERZ, Michael, Wiard; 5289 Crossbridge Drive, West Chester, OH 45069 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>

(54) Title: DIHYDROBENZOFURAN AND RELATED COMPOUNDS USEFUL AS ANTI-INFLAMMATORY AGENTS**(57) Abstract**

A compound having structure (I) wherein (a) n is from 1 to about 3; (b) X is selected from the group consisting of O, S, SO, or SO₂; (c) Y is independently hydrogen or straight, branched or cyclic alkyl having from 1 to about 4 carbon atoms, or the Y's are bonded together to form an alkanyl ring having from 3 to about 7 atoms; (d) Z is hydrogen or straight, branched or cyclic alkyl having from 3 to about 10 atoms other than hydrogen; (e) W is O or S; and (f) R₁, R₂ and R₃ are independently hydrogen, straight, branched or cyclic alkyl having from 1 to 10 carbon atoms, aryl, heterocyclyl, heteroaryl, hydroxy, or alkoxy; or R₁, R₂ or R₃ may be bonded to form one or more rings each ring having from 3 to about 7 atoms wherein from 1 to 3 atoms may be heteroatoms. Pharmaceutical compositions comprising such compounds, and methods of treating inflammation or pain using such compounds.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DIHYDROBENZOFURAN AND RELATED COMPOUNDS
USEFUL AS ANTI-INFLAMMATORY AGENTS

5

TECHNICAL FIELD

The subject invention relates to nonsteroidal anti-inflammatory drugs,
10 particularly to substituted dihydrobenzofuran and related compounds.

BACKGROUND OF THE INVENTION

Certain dihydrobenzofuran compounds and other compounds structurally
related thereto have been found to have significant disease altering activities.
Such compounds, processes for making them, and uses for them are disclosed in
15 the following references: U.S. Patent No. 4,670,457 issued to Doria, Romeo &
Corno on June 2, 1987; U.S. Patent No. 4,849,428 issued to Dobson, Loomans,
Matthews & Miller on July 18, 1989; Japanese Patent Publication No. 53-005178 of
Yoshitomi Pharm. Ind. KK published January 1, 1978; Hammond, M. L., I. E.
Kopka, R. A. Zambias, C. G. Caldwell, J. Boger, F. Baker, T. Bach, S. Luell & D. E.
20 MacIntyre, "2,3-Dihydro-5-benzofuranols as Antioxidant-Based Inhibitors of
Leukotriene Biosynthesis", J. Med. Chem., Vol. 32 (1989), pp. 1006-1020; Ortiz de
Montellano, P. R. & M. A. Correia, "Suicidal Destruction of Cytochrome P-450
during Oxidative Drug Metabolism", Ann. Rev. Pharmacol. Toxicol., Vol. 23 (1983),
pp. 481-503; Chakrabarti, J.K., R.J. Eggleton, P.T. Gallagher, J. Harvey, T.A.
25 Hicks, E.A. Kitchen, and C.W. Smith, "5-Acyl-3-substituted-benzofuran-2(3H)-ones
as Potential Anti-inflammatory Agents", J. Med. Chem., Vol. 30 (1987), pp. 1663-
1668.

It is an object of the subject invention to provide compounds which have
effective anti-inflammatory, analgesic and/or anti-oxidant activity.


30 It is a further object of the subject invention to provide such compounds
which cause few adverse side effects.

It is also an object of the subject invention to provide methods for treating
inflammation and/or pain using the subject compounds.

SUMMARY OF THE INVENTION

35 The subject invention compounds having the structure:

2



wherein

- (a) n is from 1 to about 3;
- (b) X is selected from the group consisting of O, S, SO, or SO₂;
- 5 (c) Y is independently hydrogen or straight, branched or cyclic alkyl having from 1 to about 4 carbon atoms, or the Y's are bonded together to form an alkanyl ring having from 3 to about 7 atoms;
- (d) Z is hydrogen or straight, branched or cyclic alkyl having from 3 to about 10 atoms other than hydrogen;
- 10 (e) W is O or S; and
- (f) R₁, R₂ and R₃ are independently hydrogen, straight, branched or cyclic alkyl having from one to 10 carbon atoms, aryl, heterocyclyl, heteroaryl, hydroxy, or alkoxy; or R₁, R₂ and R₃ may be bonded to form one or more rings each ring having from 3 to about 7 atoms wherein one to three atoms may be heteroatoms.
- 15

DETAILED DESCRIPTION OF THE INVENTION

As used herein, unless otherwise indicated, "alkyl" or "alkanyl" means a straight, branched or cyclic hydrocarbon chain, saturated or unsaturated, unsubstituted or substituted. Preferred alkyl are C₁-C₁₀; more preferred are C₁-C₈; especially preferred are C₁-C₄. Preferred alkyl are straight chain. Preferred branched alkyl have one or two branches, preferably one branch. Preferred cyclic alkyl are monocyclic or are a straight chain with a monocyclic terminus. Preferred alkyl are saturated. Unsaturated alkyl have one or more double bonds or/and one or more triple bonds. Preferred unsaturated alkyl have one or two double bonds or one triple bond, more preferably one double bond. Preferred alkyl are unsubstituted. Preferred substituted alkyl are mono-, di-, or trisubstituted, more preferably monosubstituted. Preferred alkyl substituents include halo, hydroxy, oxo, alkoxy (e.g., methoxy, ethoxy, propoxy, butoxy, pentoxy), aryloxy (e.g., phenoxy, chlorophenoxy, tolyloxy, methoxyphenoxy, benzyloxyphenoxy, alkyloxycarbonylphenoxy, acyloxyphenoxy), acyloxy (e.g., propionyloxy, benzoyloxy, acetoxo), carbamoyloxy, carboxy, mercapto, alkylthio, acylthio, arylthio (e.g., phenylthio, chlorophenylthio, alkylphenylthio, alkoxyphenylthio, benzylthio, alkyloxycarbonylphenylthio), aryl (e.g., phenyl, tolyl, alkyloxyphenyl,

alkyloxycarbonylphenyl, halophenyl), heterocyclyl, heteroaryl, amino (e.g., amino, mono- and di- C₁-C₃ alkylamino, methylphenylamino, methylbenzylamino), C₁-C₃ alkylamido, ureido, N'-alkylureido, N',N'-dialkylureido, N',N',N'-trialkylureido, guanidino, N'-alkylguanidino, N',N''-dialkylguanidino, or alkoxy carbonyl.

5 As used herein, "alkoxy" means -O-alkyl.

As used herein, "aryl" means a moiety having an unsubstituted or substituted aromatic ring having 6 to about 10 carbon atoms. Preferred aryl are phenyl and naphthyl; most preferred aryl is phenyl. Preferred aryl are unsubstituted. Preferred substituted aryl are mono-, di-, or trisubstituted, more preferably monosubstituted. Preferred aryl substituents include hydroxy, mercapto, halo, methyl, ethyl and propyl.

As used herein, "heterocyclyl" means a moiety having a saturated or unsaturated non-aromatic ring having from 3 to about 8 ring atoms, including from 2 to about 6 carbon atoms and from 1 to about 4 heteroatoms selected from O, S, and N. Preferred heterocycles are saturated. Preferred heterocycles have 5 or 6 atoms in the ring including 1 or 2 heteroatoms in the ring, also preferably 1 heteroatom in the ring. Specific preferred heterocycles include piperidinyl, tetrahydrothienyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrofuranyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, azepinyl, oxepinyl, thiepinyl, triazolidinyl, tetrazolidinyl. Heterocycles are unsubstituted or substituted, preferably unsubstituted. Preferred substituted heterocycles are mono-, di-, or trisubstituted, more preferably monosubstituted. Preferred heterocycle substituents include alkyl, halo, hydroxy, alkoxy, acyloxy, carboxy, carbamyloxy, thio, amino, amido, ureido, guanidino, thiocarbamamido, thioureido.


As used herein, "heteroaryl" means a moiety having an aromatic ring of 5 or 6 atoms including from 1 to 5 carbon atoms and from 1 to 4 heteroatoms selected from O, S, and N. Preferred heteroaryl groups include 1 to 3 heteroatoms in the ring, also preferably 1 or 2 heteroatoms in the ring. Specific preferred heterocycles include furyl, thienyl, pyrrolyl either unsubstituted or alkyl substituted on nitrogen, thiazolyl, oxazolyl, 5-imidazolyl either unsubstituted or alkyl-substituted on nitrogen, isoxazolyl, isothiazolyl, pyrazolyl unsubstituted or alkyl-substituted on nitrogen, oxdiazolyl, thiadiazolyl, triazolyl, tetrazolyl, pyridyl, pyrimidinyl, pyrazinyl. Fused heteroaryls include imidazothiazolinyl, imidazopyridinyl, imidazoimidazolinyl, indolyl, quinolyl, isoquinolyl. Heteroaryl groups are unsubstituted or substituted, preferably unsubstituted. Preferred substituted heteroaryls are mono-, di-, or trisubstituted, more preferably monosubstituted. Preferred heteroaryl substituents

include alkyl, halo, hydroxy, alkoxy, thio, nitro, amino, nitro, amido, ureido, guanidino, thiocarbamamido, thioureido.

As used herein, "halo" means fluoro, chloro, bromo or iodo. Preferred halo are fluoro, chloro and bromo; more preferred are chloro and bromo, especially chloro.

Compounds

The subject invention involves compounds having the following structure:



wherein

- (a) n is from 1 to about 3;
- (b) X is selected from the group consisting of O, S, SO, or SO₂;
- (c) Y is independently hydrogen or straight, branched or cyclic alkyl having from 1 to about 3 carbon atoms, or the Y's are bonded together to form an alkanyl ring having from 3 to about 7 atoms other than hydrogen;
- (d) Z is hydrogen or straight, branched or cyclic alkyl having from 3 to about 10 atoms other than hydrogen;
- (e) W is O or S; and
- (f) R₁, R₂ and R₃ are independently hydrogen, straight, branched or cyclic alkyl having from 1 to about 10 carbon atoms, aryl, heterocyclyl, heteroaryl, hydroxy, or alkoxy; or R₁, R₂ and R₃ may be bonded together to form one or more rings each ring having from 3 to about 7 atoms wherein one to three atoms may be heteroatoms.


In the above structure, each Y is independently selected from hydrogen, unsubstituted straight or branched alkanyl having from 1 to about 4 carbon atoms, and cyclic alkyl having about 3 carbon atoms, (e.g., cyclopropyl), or the Y's are bonded together to form an unsubstituted cyclic alkanyl ring having from 3 to about 7 carbon atoms in the ring. Each Y is preferably hydrogen, methyl, ethyl or cyclopropyl; more preferably hydrogen or methyl; most preferably methyl. Preferably both Y's are the same. When the Y's are bonded together to form a cyclic ring, the ring is preferably cyclopropyl, cyclobutyl or cyclopentyl, more preferably cyclopropyl.

In the above structure, Z is selected from unsubstituted branched or cyclic alkyl, and unsubstituted or alkanyl-substituted phenyl, or benzyl, Z having from 3 to about 10 atoms other than hydrogen. Z is preferably saturated.

Z is preferably branched alkanyl having from about 3 to about 10 carbon atoms, more preferably from about 4 to about 6 carbon atoms. Z is preferably branched alkanyl having 2 or more branches, more preferably 2 branches. Preferred branched alkanyl Z include t-butyl, neopentyl, isopropyl; most preferred is t-butyl. Preferred cyclic alkanyl Z include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl. Also preferred cyclic alkanyl Z include methyl or ethyl with a terminal cyclopropyl, cyclobutyl or cyclopentyl, especially cyclopropylmethyl or cyclopropylethyl. Also preferred Z is unsubstituted phenyl or benzyl.

In the above structure, R₁, R₂ and R₃ are independently hydrogen, straight, branched or cyclic alkyl having from one to 10 carbon atoms. R₁ and R₂ and/or R₂ and R₃ may be bonded together to form a cyclic alkanyl ring having from about 3 to about 7 carbon atoms in the ring where 1 to 3 atoms may be heteroatoms selected from O, N, or S. Other preferred R groups include aryl, heterocyclyl, heteroaryl, hydroxy, or alkoxy.

Preferred compounds of the subject invention are included in the following table:



20

Compound
No.

R₃

25	1	methyl
	2	ethyl
	3	propyl

In order to determine and assess pharmacological activity, testing of the subject compounds in animals is carried out using various assays known to those skilled in the art. The anti-inflammatory activity of the subject compounds can be conveniently demonstrated using an assay designed to test the ability of the subject compounds to antagonize the local edema which is characteristic of the

30

inflammatory response. Examples of such known tests include the rat carrageenan edema test, the oxazolone-induced inflamed mouse ear test, and the mouse arachadonic acid-induced inflamed ear test. Analgesic activity may be tested in art-known models such as the phenylbenzoquinone-induced writhing test in mice, and the Randall & Selitto test in rats. Another useful art-known test is the rat adjuvant arthritis test which is a useful model for assessing anti-inflammatory activity, anti-arthritic and anti-resorptive activity in a chronic, rather than an acute, model.

These and other appropriate tests for pharmacological activity are disclosed and/or referred to in U.S. Patent No. 4,130,666 issued to Moore on December 19, 1978; U.S. Patent No. 4,431,656 issued February 14, 1984 to Katsumi, et al.; U.S. Patent No. 4,440,784 issued to Katsumi, et al. on April 3, 1984; Japanese Patent Application 85/54315 of Katsumi, et al., published March 28, 1985; European Patent Application No. 0,059,090 of Yamanuchi Pharmaceutical Company Ltd., published September 1, 1982; Opas, E.V., R.J. Bonney & J. L. Humes, "Prostaglandin and Leukotriene Synthesis in Mouse Ears Inflamed by Arachadonic Acid", The Journal of Investigative Dermatology, Vol. 84, No. 4 (1985), pp. 253-256; Swingle, K. F., R. L. Bell & G. G. I. Moore, "Anti-inflammatory Activity of Antioxidants", Anti-inflammatory and Antirheumatic Drugs, Vol. III, Chapter 4, K. D. Rainsford, ed., CRC Press, Inc., (1985), pp. 105-126; Adamkiewicz, V. W., W. B. Rice & J. D. McColl, "Antiphlogistic Effect of Trypsin in Normal and in Adrenalectomized Rats", Canadian Journal of Biochemistry & Physiology, Vol. 33 (1955), pp. 332-339; Sellye, H., "Further Studies Concerning the Participation of the Adrenal Cortex in the Pathogenesis of Arthritis", British Medical Journal, Vol. 2 (1949), pp. 1129-1135; and Winter, C.A., E. A. Risley & G. W. Nuss, "Carrageenan-Induced Edema in Hind Paw of the Rats as an Assay for Antiinflammatory Drugs" Proceedings of Society of Experimental Biology and Medicine, Vol. 111 (1962), pp. 544-547; Otterness, I., & M. L. Bliven, "Laboratory Methods for Testing Nonsteroidal Antiinflammatory Drugs", Nonsteroidal Antiinflammatory Drugs, Chapter 3, J. G. Lombardino, ed., John Wiley & Sons, Inc. (1985), pp. 111-252. Hitchens, J. T., S. Goldstein, L. Shemano & J. M. Beiler, "Analgesic Effects of Irritants in Three Models of Experimentally-Induced Pain", Arch. Int. Pharmacodyn., Vol. 169, No. 2 (1967) pp. 384-393; Milne, G. M. & T. M. Twomey, "The Analgetic Properties of Piroxicam in Animals and Correlation with Experimentally Determined Plasma Levels", Agents and Actions, Vol. 10, No. 1/2 (1980), pp. 31-37; Randall, L. O. & J. J. Selitto, "A Method for Measurement of Analgesic Activity on Inflamed Tissue", Arch. Int. Pharmacodyn., Vol. 111, No. 4


(1957), pp. 409-419; Winter, C. A. & L. Faltaker, "Nociceptive Thresholds as Affected by Parenteral Administration of Irritants and of Various Antinociceptive Drugs", J. Pharmacol. Exp. Ther., Vol. 148, No. 3 (1965), pp. 373-379; the disclosure of all these references are incorporated herein by reference.

5 Many anti-inflammatory drugs, particularly non-steroidal anti-inflammatory drugs (NSAIDs) cause undesirable gastrointestinal side effects, especially when dosed perorally; such side effects may include ulcers and erosions. These side effects, which are often asymptomatic, can become serious enough to require hospitalization and can even be lethal. Compounds of the subject invention
10 generally cause fewer such gastrointestinal side effects compared to other NSAIDs. Some compounds of the subject invention are even gastroprotective, protecting the stomach and intestines from ulcers and erosions, particularly those caused by ethanol or other NSAIDs.


 Certain NSAIDs, when dosed systematically, cause an undesirable increase
15 in systemic levels of certain liver enzymes. Compounds of the subject invention generally cause little or no liver enzyme side effects.

 Compounds useful in the subject invention can be made using the following general reaction schemes:

20 The starting substituted benzene compound can be nitrated with nitric acid in sulfuric acid and the resulting nitro compound can be catalytically reduced to the corresponding aniline. This aniline is the key intermediate to ureas with different degrees of substitution. Reaction with an isocyanate or thioisocyanate gives the N, N'-disubstituted urea or thiourea. The same aniline compound can be reacted with
25 phosgene or thiophosgene to give the corresponding isocyanate or thioisocyanate which can be reacted with an amine to give the N, N', N'-trisubstituted urea or thiourea. Reductive amination using an aldehyde with the aniline intermediate gives the substituted aniline which is then reacted with an isocyanate or thioisocyanate to give the N, N'-disubstituted urea or thiourea.



The nitrated benzene can be reduced with zinc dust to provide the hydroxylamine which is then reacted with an isocyanate or thioisocyanate to provide the N-hydroxy-N'-substituted urea or thiourea.



Synthesis Examples

The following non-limiting examples provide further information regarding synthesis of the subject compounds.

Example 1N-(7-*tert*-Butyl-2,3-dihydro-3,3-dimethyl-5-benzofuranyl)-N'-methyl urea:Step 1: 7-*tert*-Butyl-2,3-dihydro-3,3-dimethyl-5-nitro-benzofuran

5

To a solution of 7-*tert*-butyl-2,3-dihydro-3,3-dimethylbenzofuran (10.0 g 49.0 mmol) in glacial acetic acid (80 mL) is added dropwise 70% HNO₃ (4.0 mL, 5.7 g, 63.7 mmol). The reaction mixture darkens to a deep green color over the course of the addition. The reaction is monitored by TLC (2% EtOAc / hexanes), and is allowed to stir at 22 °C for 4 h. Then the reaction mixture is partitioned between Et₂O (100 mL) and H₂O (2 x 100 mL). The ethereal layer was washed with saturated aqueous Na₂CO₃ (50 mL), dried (MgSO₄), filtered and evaporated to a red solid (9.12 g). Crystallization from hexane provides the title compound as orange prisms (2.26 g, 18.5%). A second crop yields slightly less pure material (4.93 g, 40.3%) which was used without further purification, mp = 93-94 °C.

15

5-Amino-7-*tert*-butyl-2,3-dihydro-3,3-dimethylbenzofuran:

A suspension of 7-*tert*-butyl-2,3-dihydro-3,3-dimethyl-5-nitro-benzofuran (2.2 g, 8.8 mmol) and 10% palladium on carbon (200 mg) in absolute EtOH (60 mL) is hydrogenated at 40 psi of H₂ pressure on a Parr hydrogenation apparatus for 3 h at 22 °C. The reaction is judged complete by TLC (hexane:EtOAc, 19:1; visualized with Dragendorff reagent). The reaction mixture is filtered through Celite and evaporated to yield the title compound as a purple solid (1.7 g, 88%), which is used without further purification.

25

Step 3: N-(7-*tert*-Butyl-2,3-dihydro-3,3-dimethyl-5-benzofuranyl)-N'-methyl urea

Methylisocyanate (0.14 mL, 2.28 mmol) is added dropwise to a solution of 5-amino-7-*tert*-butyl-2,3-dihydro-3,3-dimethylbenzofuran (500 mg, 2.28 mmol) in Et₂O (5 mL). A solid forms over 0.5 h and the reaction is quenched with H₂O (10 mL) after 1 h. The reaction is extracted with Et₂O (3 x10 mL) and the organic layers dried (MgSO₄) and evaporated to a tan solid (550 mg). This solid is recrystallized from EtOAc to give the product as white prisms (350 mg, 56%), mp = 193-194 °C.

30

Example 2N-(7-*tert*-Butyl-2,3-dihydro-3,3-dimethyl-5-benzofuranyl)-N'-ethyl urea

5 Ethylisocyanate (0.18 mL, 2.28 mmol) is added dropwise to a solution of 5-amino-7-*tert*-butyl-2,3-dihydro-3,3-dimethylbenzofuran (500 mg, 2.28 mmol) in Et₂O (5 mL). A solid forms after 10 min and the reaction is quenched with H₂O (10 mL) after 40 min. The reaction is extracted with Et₂O (3 x 10 mL) and the organic layers dried (MgSO₄) and evaporated to a tan solid (657 mg). This solid is
10 recrystallized from EtOAc to give the title compound as white prisms (335 mg, 51%), mp = 197-198 °C.

Example 3N-(7-*tert*-Butyl-2,3-dihydro-3,3-dimethyl-5-benzofuranyl)-N'-propyl urea

15 Propylisocyanate (0.21 mL, 2.28 mmol) is added dropwise to a solution of (7-*t*-butyl-2,3-dihydro-3,3-dimethyl-5-benzofuranyl)-amine (500 mg, 2.28 mmol) and Et₂O (5 mL). A solid forms after 10 min and the reaction is quenched with H₂O (10 mL) after 40 min. The reaction is extracted with Et₂O (3 x 10 mL) and the organic
20 layers dried (MgSO₄) and evaporated to an off-white solid (650 mg). This solid is recrystallized from EtOAc to give a white solid (270 mg, 39%), mp = 193 - 194 °C.

Example 4N-(7-*tert*-Butyl-2,3-dihydro-3,3-dimethyl-5-benzothieryl)-N'-methyl urea

25 Step 1. 2-Bromo-6-*tert*-butylthiophenol. To a solution of tetramethylethylenediamine (198.4 mmol, 30 mL) in cyclohexane (140 mL) is slowly added *n*-BuLi (198.4 mmol, 99.2 mL; 2 M solution in cyclohexane) at 23 °C. The resulting solution is cooled to 0 °C. A solution of 2-*tert*-butylthiophenol (15.0 g, 90.2 mmol) in cyclohexane (40 mL) is then added at a rate such that the
30 temperature stays below 10°C. The reaction is then stirred at 0 °C for 5 h and is then allowed to warm to 23 °C overnight. To the resulting yellow solution at 23 °C is added *sec*-BuLi (90.2 mmol, 69.4 mL of 1.3 M solution in cyclohexane) over 0.5 h. The resulting solution gradually turns orange. After 1.5 h, the orange, cloudy
35 reaction mixture is cannulated into a stirring solution of 1,2-dibromotetrafluoroethane (180.4 mmol, 21.5 mL) in THF (50 mL) over 1 h. After addition is complete, the resulting reaction mixture is quenched with 1 N HCl (80

mL), and extracted with hexanes (3 x 100 mL). The hexanes are dried (MgSO₄) and evaporated to a dark oil (29.48 g). This oil is taken up in 1 N NaOH (100 mL), and extracted with CH₂Cl₂ (3 x 50 mL). The organic phase is discarded, and the aqueous phase is acidified with 12 N HCl, and extracted with CH₂Cl₂ (3 x 100 mL).
5 The organic phase is dried (MgSO₄) and evaporated to provide the title compound as a yellow oil.

Step 2. 2-Bromo-6-*tert*-butylphenyl 2-methallyl thioether. A solution of 2-bromo-6-*tert*-butylthiophenol (12.4 g, 50.6 mmol), K₂CO₃ (8.44 g, 61.1 mmol), NaI (766 mg, 50.6 mmol), and β-methallylchloride (5.17 mL, 50.6 mmol) in acetone (250 mL) is
10 heated at reflux for 2 h. The reaction is monitored by TLC (hexanes). The reaction mixture is allowed to cool to 23 °C, and the precipitated solids are filtered off. The filtrate is evaporated to a dark yellow oil, which is taken up in hexanes (100 mL) and stirred with silica gel (10 g) for 20 min. The silica gel is filtered off and
15 discarded, and the filtrate is evaporated to yield the title compound as a light yellow oil.

Step 3. 7-*tert*-Butyl-2,3-dihydro-3,3-dimethylbenzothiophene. A solution of (2-bromo-6-*tert*-butylphenyl)-(2-methallyl)-thioether (9.00 g, 30.0 mmol), di-*iso*-propylethylamine (160 mL, 0.90 mol) and 80% aqueous hypophosphorous acid (58 g, 0.90 mol) in dioxane (450 mL) is deoxygenated by bubbling with N₂ for 0.5 h. The solution is heated to reflux, and a similarly deoxygenated solution of azo-bis-*iso*-butyrylnitrile (1.7 g, 8.8 mmol) in dioxane (5 mL) is added via syringe in 0.5 mL
20 portions in 0.5 h intervals. The reaction is monitored by TLC (hexanes). After 24 h, the reaction mixture is allowed to cool to 23 °C, and partitioned with 1 N HCl (300 mL), brine (100 mL), and Et₂O (3 x 150 mL). The combined ethereal extracts are back extracted with 1 N NaOH (3 x 50 mL) and H₂O (2 x 50 mL), and then are dried (MgSO₄) and evaporated to yield the crude product as a yellow oil. Short
25 path vacuum distillation (85 °C, 40 mm Hg) of this material provides the title compound as a faint yellow oil.
30

Step 4. 7-*tert*-Butyl-2,3-dihydro-3,3-dimethyl-5-nitro-benzothiophene. To a solution of 7-*tert*-butyl-2,3-dihydro-3,3-dimethylbenzothiophene (4.0 g, 18.2 mmol) in glacial acetic acid (30 mL) is added dropwise 70% HNO₃ (2.0 mL, 2.8 g, 31.8 mmol). The
35 reaction mixture darkens to a deep green color over the course of the addition. The reaction is monitored by TLC (2% EtOAc / hexanes). Then the reaction mixture is partitioned between Et₂O (50 mL) and H₂O (2 x 50 mL). The ethereal

layer is washed with saturated aqueous Na_2CO_3 (50 mL), dried (MgSO_4), filtered and evaporated to a red solid. Crystallization from hexane provides the title compound as orange prisms. A second crop yielded slightly less pure material which is combined with the former and used without further purification.

5

Step 5. 5-Amino-7-*tert*-butyl-2,3-dihydro-3,3-dimethylbenzofuran. A suspension of 7-*tert*-butyl-2,3-dihydro-3,3-dimethyl-5-nitro-benzothiophene (2.6 g, 8.8 mmol) in absolute EtOH (60 mL) is hydrogenated at 40 psi of H_2 pressure on a Parr hydrogenation apparatus for 3 h at 22 °C. The reaction is judged complete by TLC (hexane:EtOAc, 19:1; visualized with Dragendorff reagent). The reaction mixture is filtered through Celite and evaporated to yield the title compound as a purple solid, which is used without further purification.

10

Step 6. N-(7-*tert*-Butyl-2,3-dihydro-3,3-dimethyl-5-benzothiophenyl)-N'-methyl urea. Methylisocyanate (0.14 mL, 2.28 mmol) is added dropwise to a solution of 5-amino-7-*tert*-butyl-2,3-dihydro-3,3-dimethylbenzothiophene (550 mg, 2.33 mmol) in Et_2O (5 mL). A solid gradually forms and the reaction is quenched with H_2O (10 mL). The reaction is extracted with Et_2O (3 x 10 mL) and the organic layers dried (MgSO_4) and evaporated to a tan solid. This solid is recrystallized from EtOAc to give the product as white prisms, mp > 150 °C.

20

Compositions

Compositions of the subject invention comprise a safe and effective amount of the subject compounds, and a pharmaceutically-acceptable carrier. As used herein, "safe and effective amount" means an amount of a compound sufficient to significantly induce a positive modification in the condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgment. A safe and effective amount of a compound will vary with the particular condition being treated, the age and physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, the particular pharmaceutically-acceptable carrier utilized, and like factors within the knowledge and expertise of the attending physician.

25

30

Compositions of the subject invention preferably comprise from about 0.1% to about 99.9% by weight of a compound, more preferably from about 20% to about 80%, and most preferably from about 40% to about 70%.

35

In addition to the compound, the compositions of the subject invention contain a pharmaceutically-acceptable carrier. The term "pharmaceutically-

acceptable carrier", as used herein, means one or more compatible solid or liquid filler diluents or encapsulating substances which are suitable for administration to a human or lower animal. The term "compatible", as used herein, means that the components of the composition are capable of being commingled with the subject compound, and with each other, in a manner such that there is no interaction which would substantially reduce the pharmaceutical efficacy of the composition under ordinary use situations. Pharmaceutically-acceptable carriers must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the human or lower animal being treated.

Some examples of substances which can serve as pharmaceutically-acceptable carriers or components thereof are sugars, such as lactose, glucose and sucrose; starches, such as cornstarch and potato starch; cellulose and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose, cellulose acetate; powdered tragacanth; malt; gelatin; talc; solid lubricants, such as stearic acid, magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerin, sorbitol, mannitol, and polyethylene glycol; alginic acid; emulsifiers, such as the Tweens®; wetting agents such as sodium lauryl sulfate; coloring agents; flavoring agents, excipients; tableting agents; stabilizers; antioxidants; preservatives; pyrogen-free water; isotonic saline; and phosphate buffer solutions.

The choice of a pharmaceutically-acceptable carrier to be used in conjunction with a subject compound is basically determined by the way the compound is to be administered.

If the subject compound is to be injected, it is preferably injected non-intravenously; the preferred pharmaceutically-acceptable carrier is sterile, physiological saline, with blood compatible suspending agent, the pH of which has been adjusted to about 7.4. Such injectable compositions preferably comprise from about 1% to about 50% of the subject compound, more preferably from about 5% to about 25%, also preferably from about 10 mg to about 600 mg of the subject compound per dose.

Suitable pharmaceutically-acceptable carriers for topical application include those suited for use in lotions, creams, gels and the like. Topical compositions preferably contain from about 1% to about 50% of an emollient, more preferably from about 5% to about 25% of an emollient. Such topical compositions preferably comprise from about 0.1% to about 50%, of the subject compound, more

preferably from about 0.5% to about 10%, also preferably from about 5 mg to about 3500 mg per dose.

The preferred mode of administering the subject compound is perorally. The preferred unit dosage form is therefore tablets, capsules and the like, comprising a safe and effective amount of the compound, which is preferably from about 5 mg to about 3500 mg, more preferably from about 10 mg to about 1000 mg, and most preferably from about 25 mg to about 600 mg. The pharmaceutically-acceptable carriers suitable for the preparation of unit dosage forms for oral administration are well-known in the art. Their selection will depend on secondary considerations like taste, cost, and shelf stability, which are not critical for the purposes of the subject invention, and can be made without difficulty by a person skilled in the art.

Many of the subject compounds are hydrophobic. If it is desired to provide an aqueous-based composition or a composition soluble in or miscible with aqueous media, a solubilizing agent may be included in the composition. Non-limiting examples of such solubilizing agents include polyethylene glycol, propylene glycol, ethanol, and polyoxyethylene (35) castor oil.

Particularly preferred oral composition carriers suitable for compositions of the subject invention are disclosed in U.S. Patent Nos. 5,189,066 of Kelm & Bruns, issued February 23, 1993, entitled "Pharmaceutical Compositions of Tebufelone", and 5,281,420 of Kelm & Dobrozsi, issued January 25, 1994, entitled "Solid Dispersion Compositions of Tebufelone", hereby incorporated herein by reference.

Methods

Another aspect of the subject invention is methods for treating or preventing diseases characterized by inflammation by administering a safe and effective amount of a subject compound to a human or lower animal in need of such treatment. The term "diseases characterized by inflammation", as used herein, means conditions which are known to involve inflammation, and may include conditions such as arthritis (e.g., rheumatoid arthritis, osteoarthritis, psoriatic arthritis, juvenile arthritis, Reiter's syndrome, infectious arthritis, and ankylosing spondylitis, systemic lupus, erythematosis and gout), as well as the presence of inflammation whether or not it is associated with an identifiable disease. Diseases characterized by inflammation further may include inflammation in the oral cavity (e.g., inflammation associated with gingivitis or periodontal disease); inflammation in the gastrointestinal tract, (e.g., inflammation associated with ulcers and irritable bowel disease); inflammation associated with dermatological diseases (e.g., psoriasis, acne, and other skin inflammation); inflammation associated with the

respiratory tract (e.g., asthma, bronchitis, and allergies); and inflammation in the central nervous system (e.g., Alzheimer's disease).

Another aspect of the subject invention is methods for treating or preventing pain by administering a safe and effective amount of a subject compound to a human or lower animal in need of such treatment. Pain which can be treated or prevented by administering the subject compounds may include peripheral pain, menstrual pain, dental pain, and lower back pain.

Another aspect of the subject invention is methods for preventing oxidative damage at inflammatory sites by administering a safe and effective amount of a subject compound to a human or lower animal in need of such treatment. While not limited to a particular mechanism, it is believed that the subject compounds inhibit leukotriene synthesis, thereby decreasing neutrophil accumulation at an inflammatory site.

Another aspect of the subject invention is methods for treating or preventing gastric or duodenal ulcers or erosions by administering a safe and effective amount of a subject compound to a human or lower animal in need of such treatment. In particular, such ulcers or erosions caused by ethanol or non-steroidal antiinflammatory drugs (NSAIDs) can be treated and/or prevented by administration of preferred subject compounds.

Appropriate tests for determining the gastrointestinal safety or gastroprotective or gastric healing properties of the subject compounds are known.

Methods for determining acute gastrointestinal safety are disclosed and/or referred to in the following references: Unangst, P.C., G.P. Shrum, D.T. Connor, R.D. Dyer, and D.J. Schrier, "Novel 1,2,4-Oxadiazoles and 1,2,4-Thiadiazoles as Dual 5-Lipoxygenase and Cyclooxygenase Inhibitors", J. Med. Chem., Vol. 35 (1992), pp. 3691-3698; and Segawa, Y., O. Ohya, T. Abe, T. Omata, et al., "Anti-inflammatory, Analgesic, and Antipyretic Effects and Gastrointestinal Toxicity of the New Anti-inflammatory Drug N-{3-[3-(piperidinylmethyl)phenoxy] propyl}-carbamoylmethylthio]ethyl 1-(p-chlorobenzoyl) 5-Methoxy-2methyl-3-indolylacetate", Arzneim.-Forsch./Drug Res., Vol. 42 (1992), pp. 954-992. In the methods disclosed therein, stomachs of the animals are typically examined two hours after dosing a compound. Methods for determining subchronic gastrointestinal safety are disclosed and/or referred to in the following references: Melarange, R., C. Gentry, et al., "Anti-inflammatory and Gastrointestinal Effects of Nabumetone or Its Active Metabolite, 6-Methoxy-2-naphthylacetic Acid (6MNA)", Dig. Dis. Sci., Vol. 37 (1992), pp. 1847-1852; and Wong, S., S.J. Lee, et al.,

"Antiarthritic Profile of BF-389 - A Novel Anti-inflammatory Agent With Low Ulcerogenic Liability", Agents Actions, Vol. 37 (1992), pp. 90-91.

Methods for determining acute gastroprotection are disclosed and/or referred to in the following reference: Playford, R.J., D.A. Versey, S. Haldane, M.R. Alison, and J. Calan, "Dose-dependent Effects of Fentanyl on Indomethacin-induced Gastric Damage", Digestion, Vol. 49 (1991), pp. 198-203. In the method disclosed therein, female Lewis rats (130-175 g) are dosed perorally with the subject compound (40 mg/kg b.i.d.) or vehicle at 2 hours and immediately before administration of a gastric damaging dose of indomethacin. The rats are sacrificed 4 hours later by CO₂ asphyxiation. Gastric corpus damage (millimeters of hemorrhagic lesions) is measured by digitized imaging.

The preferred mode of administration of the subject compounds is peroral, but other known methods of administration are contemplated as well, e.g., dermatomucosally (for example, dermally, rectally and the like), and parenterally (for example, by subcutaneous injection, intramuscular injection, intraarticular injection, intravenous injection and the like). Ocular administration and inhalation are also included. Thus specific modes of administration include, without limitation, peroral, transdermal, mucosal, sublingual, intranasal, intramuscular, intravenous, intraperitoneal, subcutaneous, and topical administration.

Preferred doses of the subject compounds range from about 0.2 mg/kg to about 70 mg/kg, more preferably from about 0.5 mg/kg to about 12 mg/kg. Preferred injectable doses comprise from about 0.1 mg/kg to about 10 mg/kg of the subject compound. Preferred topical doses comprise from about 1 mg/cm² to about 200 mg/cm² of the subject compound applied to the skin surface. Preferred peroral doses comprise from about 0.5 mg/kg to about 50 mg/kg, more preferably from about 1 mg/kg to about 20 mg/kg, more preferably still from about 2 mg/kg to about 10 mg/kg, of the subject compound. Such doses are preferably administered from about once to about six times daily, more preferably from about twice to about four times daily. Such daily doses are preferably administered for at least one week, also preferably for at least two weeks, also preferably at least one month, also preferably for at least 2 months, also preferably for at least 6 months, 1 year, 2 years, or more.

Compositions and Method Examples

The following non-limiting examples illustrate the subject invention.

Example A

Pharmaceutical compositions in the form of tablets are prepared by conventional methods, such as mixing and direct compaction, formulated as follows:

	<u>Ingredient</u>	<u>Quantity (mg per tablet)</u>
5	Compound 1	200
	Microcrystalline Cellulose	100
	Sodium Starch Glycollate	30
	Magnesium Stearate	3

When administered orally two times daily, the above composition significantly reduces the inflammation in a patient suffering from rheumatoid arthritis. A significant benefit is also achieved by twice daily administration of this composition to a patient suffering from osteoarthritis.

Example B

A pharmaceutical composition in capsule form is prepared by conventional methods, formulated as follows:

	<u>Ingredient</u>	<u>Quantity (mg per capsule)</u>
15	Compound 2	200
	Lactose	To fill to volume of capsule

The above capsule administered orally once a day substantially reduces the symptomology of a patient afflicted with rheumatoid arthritis or osteoarthritis.

Example C

A pharmaceutical composition in liquid form is prepared by conventional methods, formulated as follows:

	<u>Ingredient</u>	<u>Quantity</u>
25	Compound 3	200 mg
	EtOH	4 ml
	Methyl cellulose	0.4 mg
	Distilled water	76 ml
	Tween 80	1.6 ml

50 ml of the above composition administered perorally once a day substantially reduces the symptoms of a patient afflicted with rheumatoid arthritis or osteoarthritis.

Example D

A pharmaceutical composition in liquid form is prepared by conventional methods, formulated as follows:

<u>Ingredient</u>	<u>Quantity</u>
Microcrystalline (micronoized)	200 mg

Compound 1

Avicel (microcrystalline cellulose)	50 mg
Tween 80	1.6 ml
Methyl cellulose	0.4 mg
Deionized water	80 ml

5


50 ml of the above composition administered perorally twice a day substantially reduces the symptoms of a patient afflicted with rheumatoid arthritis or osteoarthritis.

While particular embodiments of the subject invention have been described,
10 it would be obvious to those skilled in the art that various changes and modifications to the compositions disclosed herein can be made without departing from the spirit and scope of the invention. It is intended to cover, in the appended claims, all such modifications that are within the scope of this invention.

WHAT IS CLAIMED IS:

15

1. A compound having the structure:



wherein

- (a) n is from 1 to about 3;
 - 5 (b) X is selected from the group consisting of O, S, SO, or SO₂;
 - (c) Y is independently hydrogen or straight, branched or cyclic alkyl having from 1 to about 3 carbon atoms, or the Y's are bonded together to form an alkanyl ring having from 3 to about 7 atoms;
 - (d) Z is hydrogen or straight, branched or cyclic alkyl having from 3 to
10 about 10 atoms other than hydrogen;
 - (e) W is O or S; and
 - (f) R₁, R₂ and R₃ are independently hydrogen, straight, branched or
15 cyclic alkyl having from 1 to about 10 carbon atoms, aryl, heterocyclyl, heteroaryl, hydroxy, or alkoxy; or R₁, R₂ and R₃ may be bonded to form one or more rings, each ring having from 3 to about 7 atoms wherein one to three atoms may be heteroatoms.
2. The compound of Claim 1 wherein X is oxygen or sulphur, and R₁ and R₂ are hydrogen or methyl.
 3. The compound of Claim 1 wherein each Y is independently selected from the group consisting of hydrogen, methyl and ethyl; and Z is selected from the group consisting of C₄-C₆ branched alkanyl having 2 branches, unsubstituted C₃-C₆ cycloalkanyl.
 4. The compound of Claim 3 wherein X is oxygen, both Y are methyl, and Z is t-butyl.
 5. The compound of Claim 4 wherein R₃ is C₁-C₆ straight or single-branched alkyl or straight alkyl with a terminal cyclic alkyl, saturated or unsaturated with one double bond between non-terminal carbon atoms, or C₃-C₆ cycloalkanyl.

6. The compound of Claim 5 wherein X is oxygen, and R₁ and R₂ are hydrogen.
7. The compound of Claim 6 wherein n is one.
8. The compound of Claim 7 wherein R₃ is selected from the group consisting of methyl, ethyl, n-propyl, i-propyl, n-butyl, 1-methylpropyl, 2-methylpropyl, 1-methylbutyl, ethoxy, benzyl, phenethyl, cyclobutyl, cyclopentyl and cyclohexyl.
9. The compound of Claim 8 wherein W is oxygen.
10. The compound of Claim 9 wherein R₃ is selected from the group consisting of methyl, ethyl, n-propyl, n-butyl and ethoxy.
11. The compound of Claim 10 wherein R₃ is methyl, ethyl or propyl.
12. A composition comprising a compound of Claim 1 and a pharmaceutically-acceptable carrier.
13. A method of treating inflammation or pain comprising administration, to a human or lower animal in need of such treatment, of a safe and effective amount of a compound of Claim 1.
14. A method of treating arthritis comprising daily peroral administration, to a human in need of such treatment, of from about 1 mg/kg to about 20 mg/kg of a compound of Claim 1.
15. A composition comprising a compound of Claim 11 and a pharmaceutically-acceptable carrier.
16. A method of treating inflammation or pain comprising administration, to a human or lower animal in need of such treatment, of a safe and effective amount of a compound of Claim 11.

17. A method of treating arthritis comprising daily peroral administration, to a human in need of such treatment, of from about 1 mg/kg to about 20 mg/kg of a compound of Claim 11.

INTERNATIONAL SEARCH REPORT

Int. national Application No

PCT/US 97/01308

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07D307/79 C07D333/54 A61K31/34		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 73, no. 1, 6 July 1970 Columbus, Ohio, US; abstract no. 3748t, LETTIERI ET AL.: "Substituted chroman-6-ylureas and thioureas" XP002031222 &J.Med.Chem. 1970,13(3),584-585 see abstract ---	1,2
X	US 3 963 717 A (COOKE GEORGE A ET AL) 15 June 1976 see formula III columns 4-5 --- -/-	1,2
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
<div style="display: flex;"> <div style="flex: 1;"> <p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">20 May 1997</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">20.06.97</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Steendijk, M</div>

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/01308

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Section Ch, Week 8530 Derwent Publications Ltd., London, GB; Class E23, AN 85-180845 XP002031224 & JP 60 108 846 A (KONISHIROKU PHOTO IND CO LTD) , 14 June 1985 see abstract</p>	1
X	<p>DE 35 37 884 A (FMC CORP) 30 April 1986 see examples page 28</p>	1,2
X	<p>CHEMICAL ABSTRACTS, vol. 119, no. 13, 27 September 1993 Columbus, Ohio, US; abstract no. 139079b, TAKEUCHI ET AL.: "Preparation of (pyrrolidinoethyl)urea derivatives as analgesics" XP002031223 & W09309011 see abstract</p>	1,2,12
X	<p>EP 0 538 099 A (RHONE POULENC RORER SA) 21 April 1993 see claim1 and example in claim 5</p>	1,2,12
A	<p>EP 0 322 004 A (PROCTER & GAMBLE) 28 June 1989 cited in the application see the whole document</p>	1-17
P,X	<p>WO 96 07651 A (PROCTER & GAMBLE) 14 March 1996 see the whole document</p>	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/01308

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3963717 A	15-06-76	NONE	
DE 3537884 A	30-04-86	AP 36 A	06-12-88
		AT 306085 A	15-09-93
		AU 556949 B	27-11-86
		AU 4859885 A	01-05-86
		BE 903521 A	25-04-86
		BR 8505326 A	05-08-86
		CA 1265145 A	30-01-90
		CH 666029 A	30-06-88
		CS 8507615 A	15-01-87
		EG 17848 A	30-03-91
		FR 2572400 A	02-05-86
		GB 2166137 A,B	30-04-86
		JP 1482131 C	27-02-89
		JP 61106558 A	24-05-86
		JP 63026109 B	27-05-88
		LU 86134 A	11-06-86
		NL 8502913 A	16-05-86
		OA 8127 A	31-03-87
		SE 8505026 A	26-04-86
		US 4767779 A	30-08-88
EP 0538099 A	21-04-93	FR 2682381 A	16-04-93
		AT 130610 T	15-12-95
		AU 2888892 A	03-05-93
		CA 2120035 A	15-04-93
		DE 69206274 D	04-01-96
		DE 69206274 T	28-03-96
		EP 0607355 A	27-07-94
		ES 2081136 T	16-02-96
		WO 9307130 A	15-04-93
		JP 7502984 T	30-03-95
		NO 940815 A	08-03-94
		US 5563136 A	08-10-96
		ZA 9207746 A	19-04-93
EP 0322004 A	28-06-89	US 4849428 A	18-07-89
		AU 2577188 A	25-05-89
		CA 1308110 A	29-09-92

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/01308

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0322004 A		DE 3867827 A IE 61197 B IL 88365 A JP 1250372 A	27-02-92 19-10-94 27-02-94 05-10-89
WO 9607651 A	14-03-96	AU 3505995 A CA 2199128 A	27-03-96 14-03-96